

Musical rhythm spectra from Bach to Joplin obey a $1/f$ power law

Daniel J. Levitin^{a,1}, Parag Chordia^b, and Vinod Menon^{c,1}

^aDepartment of Psychology, School of Computer Science, and School of Music, McGill University, Montreal, QC, Canada H3A 1B1; ^bSchool of Music, Georgia Institute of Technology, Atlanta, GA 30332; and ^cProgram in Neurosciences, Department of Psychiatry and Behavioral Sciences and Department of Neurology and Neurological Sciences, Stanford University, Stanford, CA 94305

Edited* by Dale Purves, Duke University Medical Center, Durham, NC, and approved January 10, 2012 (received for review August 31, 2011)

Much of our enjoyment of music comes from its balance of predictability and surprise. Musical pitch fluctuations follow a $1/f$ power law that precisely achieves this balance. Musical rhythms, especially those of Western classical music, are considered highly regular and predictable, and this predictability has been hypothesized to underlie rhythm's contribution to our enjoyment of music. Are musical rhythms indeed entirely predictable and how do they vary with genre and composer? To answer this question, we analyzed the rhythm spectra of 1,788 movements from 558 compositions of Western classical music. We found that an overwhelming majority of rhythms obeyed a $1/f^\beta$ power law across 16 subgenres and 40 composers, with β ranging from ~ 0.5 – 1 . Notably, classical composers, whose compositions are known to exhibit nearly identical $1/f$ pitch spectra, demonstrated distinctive $1/f$ rhythm spectra: Beethoven's rhythms were among the most predictable, and Mozart's among the least. Our finding of the ubiquity of $1/f$ rhythm spectra in compositions spanning nearly four centuries demonstrates that, as with musical pitch, musical rhythms also exhibit a balance of predictability and surprise that could contribute in a fundamental way to our aesthetic experience of music. Although music compositions are intended to be performed, the fact that the notated rhythms follow a $1/f$ spectrum indicates that such structure is no mere artifact of performance or perception, but rather, exists within the written composition before the music is performed. Furthermore, composers systematically manipulate (consciously or otherwise) the predictability in $1/f$ rhythms to give their compositions unique identities.

musical structure | $1/f$ distributions | fractal mathematics | temporal perception

Musical behaviors—singing, dancing, and playing instruments—date back to Neanderthals (1), and have been a part of every human culture as far back as we know (2, 3). People experience great enjoyment and pleasure from music (4–6), and music theorists have argued that this enjoyment stems in part from the structural features of music, such as the generation and violation of expectations (5, 7–11). Mathematics has often been used to characterize, model, and understand music, from Schenkerian analysis (12, 13) to neural topography (14) and geometric models of tonality (15, 16). One particular mathematical relation that has received attention in music is the $1/f$ distribution, which Mandelbrot (17) termed “fractal.”

$1/f$ distributions have been found to be a key feature of a number of natural and sensory phenomena. In analyzing the frequency of several natural disasters, including earthquakes, landslides, floods, and terrestrial meteor impacts, Hsü (18) found an inverse log-log linear (fractal) relation between the frequency and the intensity of the events:

$$f = c/M^D \quad [1]$$

where f is the temporal frequency, M is a parameter indexing the intensity of the events, c is a constant of proportionality, and D is the fractal dimension. The well-known Richardson Effect

(19, 20), which began the modern study of fractals, states that measurements of natural phenomena are characterized by $1/f$ noise. $1/f$ fluctuations are also a prominent feature of human cognition. Neurons in primary visual cortex were found to exhibit higher gain, and the spike responses exhibit higher coding efficiency and information transmission rates for $1/f$ signals (21), and the fluctuations of voltage across the resting membrane of myelinated nerve fibers show a $1/f$ spectrum (22); these findings suggest that human sensory and neural systems evolved to encode certain regularities in the physical world (23, 24), in this case, those that manifest self-similarity.

The spectral power of such signals decays exponentially with frequency (f) as $[1/f]^\beta$ (where β is the spectral exponent). Although large values of β (>2) indicate greater long-range correlations, and hence, highly predictable signals, very small β s (<0.5) indicate highly unpredictable signals: An extreme example is white noise ($\beta = 0$), the structure of which is entirely unpredictable (25). On the other hand, musical pitch and loudness fluctuations are known to have an intermediate range of β values ($1 < \beta < 2$), and this range has been suggested to indicate a level of predictability that is optimal to our musical experience (26, 27).

In contrast, musical rhythms, note onsets, and durations in Western classical music are considered highly regular and predictable, so much so that the term “rhythmic” has come to be synonymous with “recurring,” “regular,” or “periodic.” Because music has a beat and is based on repetition, it has been said that “what” the next musical event will be is not always easy to guess, but “when” it is likely to happen can be easily predicted (11). In fact, this comforting predictability has been suggested to underlie rhythm's fundamental contribution to the aesthetic experience of music (7, 28, 29).

It has been previously suggested that music can be characterized by a fractal geometry (30–32), obeying a $1/f$ power law. This suggestion was demonstrated for the amplitude (26, 27) and pitch structure (33, 34) of music. Here we show that the temporal/rhythmic properties of music across a wide range of compositions also obey the $1/f$ law, or fractal relation.

We analyzed the spectral structure of musical rhythms (indexed by note onsets, and using a multitaper spectral analysis) to investigate whether these are indeed as regular and predictable as commonly claimed. Whereas previous researchers have studied only a handful of compositions by Bach, Mozart, and European folk songs (26, 27, 33, 34), we sought to increase the generalizability and importance of such findings by leveraging the power of a large database of notated musical compositions,

Author contributions: D.J.L. and V.M. designed research; D.J.L., P.C., and V.M. performed research; and D.J.L., P.C., and V.M. wrote the paper.

The authors declare no conflict of interest.

*This Direct Submission article had a prearranged editor.

¹To whom correspondence may be addressed. E-mail: daniel.levitin@mcgill.ca or menon@stanford.edu.

This article contains supporting information online at www.pnas.org/lookup/suppl/doi:10.1073/pnas.1113828109/-DCSupplemental.




Fig. 2. Musical rhythm spectra obey a $1/f$ power law. (A) Rasterized rhythm representation (Lower) showing note onsets extracted from Beethoven's Quartet Op. 18. No. 1 (score, Upper). The representation shown is schematic: actual durations were extracted from the Humdrum kern format (*Materials and Methods*). (B) (Left) The spectrum of the rhythm raster from A has power that decays linearly (in a log-scale) with frequency as $1/f$ (gray dots). The slope of the spectrum (spectral exponent or β) is 0.8. Colored segments show the sequence of durations (internote intervals). Black line represents the linear fit to the spectrum in the frequency range of 0.01 to 1 Hz (delineated by dotted vertical gray lines). Dashed line represents extrapolation of the linear fit to other frequencies. (Right) The spectrum of a sequence with the note onsets shuffled randomly, keeping durations intact. The shuffled spectrum is flat ($\beta = 0.0$). Other conventions are as shown (Left). (C) Distribution of rhythm spectral exponents pooled across genres (black) obtained by linear fits to individual pieces across the population of 1,788 pieces analyzed. Gray: spectral exponent distribution for the corresponding shuffled rhythms. Inverted triangles represent the distribution median. Dashed vertical line: $\beta = 0$.

ragtime, had among the smallest β -values, indicating the least-predictable rhythms (Fig. 3E and Fig. S1).

The β -values for compositions grouped by composer revealed a surprising trend: β -distributions varied widely across composers regardless of period or style (Fig. 4 and Table S2). Beethoven, Haydn, and Mozart demonstrated different $1/f$ rhythm spectra. Beethoven's rhythms exhibited the largest β s (and thus the most predictability), Mozart's β s were among the smallest (with the least predictability), whereas those of Haydn were intermediate between those of Mozart and Beethoven, and significantly different from either (Fig. 4 A, B, and E) [Tukey–Kramer honestly significantly different (HSD), $P < 0.05$; see *Materials and Methods*]. Monteverdi and Joplin (Table S2) exhibited nearly identical, overlapping rhythm β -distributions (Tukey–Kramer HSD, $P > 0.05$) (Fig. 4 C–E). Figs. 3E and 4E display 95% confidence intervals (CI); nonoverlapping CIs are significant at $P < 0.05$.

Discussion

Most listeners find music pleasing when it creates an optimal balance of predictability and surprise (7–9, 28, 29). Whereas such a balance was previously attributed exclusively to $1/f$ structure in musical pitch and amplitude (26, 27, 33, 34), the present results demonstrate that musical rhythms also exhibit $1/f$ spectral structure. The $1/f$ structure allows us to quantify the range of predictability, self-similarity, or fractal-like structure within which listeners find aesthetic pleasure. Critically, compositions across four centuries and several subgenres of Western classical music all demonstrated $1/f$ structure in their rhythm spectra.

A previous study analyzing classical compositions from the 18th to 20th centuries reported nearly identical $1/f$ pitch structure among composers, with a very narrow range of spectral exponents ($1.79 \leq \beta \leq 1.97$) (8). The $1/f$ pitch structure could not systematically delineate one composer's work from another. In contrast, our findings demonstrate that $1/f$ rhythm spectral exponents varied widely and systematically among composers ($0.48 \leq \beta \leq 1.05$). Even composers belonging to the same musical era (1750–1820, the Classical era), such as Beethoven, Haydn, and Mozart, demonstrated distinctive $1/f$ rhythm spectra. Conversely, Monteverdi and Joplin, composers of entirely different musical eras and composing nearly three centuries apart, exhibited similar rhythm spectra. These results suggest a heretofore underappreciated importance of rhythm and hint at its even greater role than pitch in conveying the distinctive style of composers.

Human perception is known to be sensitive to $1/f$ structure in the environment (21, 22, 25, 36, 39, 40), and electrophysiological studies have suggested a preference of sensory neural coding for $1/f$ signals (21, 39, 41). How does this relate to music? The balance of expectations both realized and violated in pleasurable music requires a certain amount of instability in the temporal structure, and such dynamical instability may best be modeled by a power law for temporal fluctuation (42).

Cognitive psychologists (43) have noted that human performance in a variety of tasks, including the production of rhythmic sequences by finger tapping (44), fluctuates over time according to $1/f$. Here, we reveal that this same structure characterizes the stage of human cognition before action: the written composition of temporal intervals that will become action plans only at a later date when performed by musicians. Thus, $1/f$ is not merely an artifact of performance, but exists in the written scores themselves. Perhaps composers can't help but produce $1/f$ rhythm spectra, perhaps musical conventions require this. From a psychological standpoint, the finding suggests that composers have internalized some of the regularities of the physical world as it interacts with biological systems (including the mind) to recreate self-similarity in works of musical art (cf. 24).

Our finding of $1/f$ rhythm structure in nearly four centuries of musical compositions may therefore be rooted in a fundamental propensity of our sensory and motor systems to both perceive and produce $1/f$ structure (44, 45).

Finally, an important methodological contribution of our study is the development of raster representations borrowed from the neuroscience literature, combined with a multitaper spectral analysis (*SI Materials and Methods*) to rigorously examine rhythm structure in music. Although pitch and loudness are musical attributes that can be represented as a continuous signal having a specific value at each instant of time (36), musical rhythms are essentially sequences of note durations (intervals of time) that, by definition, cannot be represented as a continuous signal with a specific value at each point in time. Representation of rhythmic structure as point processes allows us to examine spectral and temporal dynamics with unprecedented mathematical precision and rigor. The methods developed in our study are likely to be useful for addressing many other important questions involving rhythmicity in music and speech across cultures.

Materials and Methods

Analysis of Musical Scores. All musical scores were drawn from the Humdrum Kern database (46). Voices (instruments) and durations were extracted with custom scripts using the Humdrum Toolkit (47). Musical movements, which are self-contained sections of larger compositions, were treated as independent units (pieces). Onsets for each voice of each piece were converted into a time-aligned raster representation, providing a marker of interonset intervals (interevent durations). Onsets from different voices were merged (as shown in Fig. 2A). Rhythm spectra were then computed with the multitaper approach using the Chronux toolbox (35, 48). The multitaper approach (35) minimizes the variance (and spectral leakage) of the spectral estimate

exponent means based on the Tukey–Kramer HSD criterion. Statistical analyses were performed in Matlab. We repeated the spectral analysis for each voice extracted separately, normalizing all pieces to the same total duration, as well as including pieces of all lengths (< 200 notes), and observed results similar to those reported here. Plots of the power spectra for individual composers and genres, as well as additional details on the compositions used in this study, can be found in *SI Appendix 1* and *SI Appendix 2*.

Generation of Simulated 1/f Rhythms. To generate duration sequences that obey a power law of the form $[1/f]^\beta$ we modeled the internote intervals, or durations, (τ) as a multiplicative stochastic point process (53):

$$\tau_{k+1} = \tau_k + \gamma \tau_k^{2\mu-1} + \sigma \tau_k^\mu \epsilon_k \quad [2]$$

In this formula, γ is a relaxation factor, e represents a normally distributed, zero-mean, unit-variance, white-noise process, and σ represents a scaling factor for the SD of the white noise. The relaxation factor (γ) was varied to generate duration sequences with different spectral exponents (β). Other parameters were set to fixed values ($\mu = 0.0$, $\sigma = 0.05$). Larger values of the

relaxation factor led to sequences with longer (persistent) history effects: such sequences showed correlations over longer time windows (Fig. 1A), and exhibited 1/f spectra with larger β s (Fig. 1B). Duration sequences with $\beta = 2.0$ were generated with $\gamma = 0.025$, whereas sequences with $\beta = 1.0$ were generated with $\gamma = 0.0$. Sequences with $\beta = 0.0$ (white noise) were generated with $\gamma = 0.0$, but without the first term on the right hand side of Eq. 1, thus eliminating all history effects. Rhythm rasters were then derived from the duration sequences by placing “ticks” on the time axis such that successive ticks in time were separated by successive values in the duration sequence. All simulations were performed in Matlab.

ACKNOWLEDGMENTS. We thank Evan Balaban for assistance in developing the original idea for this study and Devarajan Sridharan for his help with generation and analysis of simulated rhythm sequences, rasterized representation of the musical rhythms from the Center for Computer Assisted Research in the Humanities database, multitaper spectral analysis, statistical analysis of rhythm spectra, and with figures and tables. This work was supported by National Science Foundation Grants BCS-0449927 (to V.M. and D.J.L.) and IIS-0855758 (to P.C.), by Natural Sciences and Engineering Research Council Grant 228175-10 and a Google Faculty Research Award (to D.J.L.).

- Mithen S (2006) *The Singing Neanderthals: The Origins of Music, Language, Mind, and Body* (Harvard Univ Press, Cambridge, MA).
- Cross I (1999) *Music, Mind and Science*, ed Yi SW (Western Music Research Institute, Seoul National University, Seoul), pp 35–40.
- Cross I (2001) Music, cognition, culture, and evolution. *Ann N Y Acad Sci* 930:28–42.
- Plato (1987) *The Republic* (Penguin, London).
- Huron D (2006) *Sweet Anticipation: Music and the Psychology of Expectation* (MIT Press, Cambridge, MA).
- Seashore CE (1967) *Psychology of Music* (Dover, Mineola, NY).
- Meyer L (1956) *Emotion and Meaning in Music* (Univ of Chicago Press, Chicago).
- Lerdahl F, Jackendoff R (1983) *A Generative Theory of Tonal Music* (MIT, Cambridge).
- Levitin DJ (2006) *This Is Your Brain on Music* (Dutton, New York).
- Narmour E (1992) *The Analysis and Cognition of Melodic Complexity: The Implication-Realization Model* (Univ of Chicago Press, Chicago).
- Levitin DJ (2010) Why music moves us. *Nature* 464:834–835.
- Forte A, Gilbert SE (1982) *Introduction to Schenkerian Analysis: Form and Content in Tonal Music* (Norton, New York).
- Schenker H (1954) *Harmony*, trans Borgese E (Univ of Chicago Press, Chicago).
- Janata P, et al. (2002) The cortical topography of tonal structures underlying Western music. *Science* 298:2167–2170.
- Callender C, Quinn I, Tymoczko D (2008) Generalized voice-leading spaces. *Science* 320:346–348.
- Tymoczko D (2006) The geometry of musical chords. *Science* 313:72–74.
- Mandelbrot BB (1977) *The Fractal Geometry of Nature* (Freeman, New York).
- Hsü KJ (1983) Actualistic catastrophism: Address of the retiring president of the international association of sedimentologists. *Sedimentology* 30(1):3–9.
- Mandelbrot B (1967) How long is the coast of Britain? Statistical self-similarity and fractional dimension. *Science* 156:636–638.
- Richardson LF (1961) The problem of contiguity: An appendix to statistic of deadly quarrels. *GenSys: Yearbk Soc Adv Gen Sys Thry* 6(139):129–187.
- Yu Y, Romero R, Lee TS (2005) Preference of sensory neural coding for 1/f signals. *Phys Rev Lett* 94:108103.
- Verveen AA, Derksen HE (1968) Fluctuation phenomena in nerve membrane. *Proc IEEE* 56:906–916.
- Shepard RN (1994) Perceptual-cognitive universals as reflections of the world. *Psychon Bull Rev* 1(1):2–28.
- Shepard RN (2001) Perceptual-cognitive universals as reflections of the world. *Behav Brain Sci* 24:581–601, discussion 652–671.
- Schmuckler MA, Gilden DL (1993) Auditory perception of fractal contours. *J Exp Psychol Hum Percept Perform* 19:641–660.
- Voss RF, Clarke J (1978) “1/f noise” in music: Music from 1/f noise. *J Acoust Soc Am* 63: 258–263.
- Voss RF, Clark J (1975) 1/f noise in music and speech. *Nature* 258:317–318.
- Bernstein L (1959) *The Joy of Music* (Simon and Schuster, New York).
- Bernstein L (1976) *The Unanswered Question: Six Talks at Harvard (Charles Eliot Norton Lectures)* (Harvard Univ Press, Cambridge, MA).
- Campbell P (1986) The music of digital computers. *Nature* 324:523–528.
- Campbell P (1987) Is there such a thing as fractal music? *Nature* 325:766.
- Schroeder MR (1987) Is there such a thing as fractal music? *Nature* 325:765–766.
- Hsü KJ, Hsü AJ (1990) Fractal geometry of music. *Proc Natl Acad Sci USA* 87:938–941.
- Hsü KJ, Hsü AJ (1991) Self-similarity of the “1/f noise” called music. *Proc Natl Acad Sci USA* 88:3507–3509.
- Mitra P, Bokil H (2008) *Observed Brain Dynamics* (Oxford Univ Press, New York).
- Boon JP, Decroly O (1995) Dynamical systems theory for music dynamics. *Chaos* 5: 501–508.
- Abrams DA, et al. (2011) Decoding temporal structure in music and speech relies on shared brain resources but elicits different fine-scale spatial patterns. *Cereb Cortex* 21:1507–1518.
- Levitin DJ, Menon V (2003) Musical structure is processed in “language” areas of the brain: A possible role for Brodmann Area 47 in temporal coherence. *Neuroimage* 20: 2142–2152.
- Patel AD, Balaban E (2000) Temporal patterns of human cortical activity reflect tone sequence structure. *Nature* 404:80–84.
- García-Lázaro JA, Ahemd B, Schnupp JW (2006) Dynamical substructure of coordinated rhythmic movements. *Curr Biol* 16:264–271.
- Wu D, Li C-Y, Yao D-Z (2009) Scale-free music of the brain. *PLoS ONE* 4:e5915.
- Bak P, Tang C, Wiesenfeld K (1987) Self-organized criticality: An explanation of the 1/f noise. *Phys Rev Lett* 59:381–384.
- Wagenmakers E-J, Farrell S, Ratcliff R (2004) Estimation and interpretation of 1/falpa noise in human cognition. *Psychon Bull Rev* 11:579–615.
- Gilden DL, Thornton T, Mallon MW (1995) 1/f noise in human cognition. *Science* 267: 1837–1839.
- Schmidt RC, Beek PJ, Treffner PJ, Turvey MT (1991) Dynamical substructure of coordinated rhythmic movements. *J Exp Psychol Hum Percept Perform* 17:635–651.
- Huron D (2010) Humdrum Kern database, <http://kern.humdrum.net>. Accessed August 15, 2010.
- Huron D (1998) Humdrum Toolkit, <http://humdrum.org/Humdrum/>. Accessed August 15, 2010.
- Mitra-Laboratory (MatLab, Cold Spring Harbor, NY). Available at <http://chronux.org/chronux/>. Accessed August 15, 2010.
- Slepian D (1978) Prolate spheroidal wave functions, Fourier analysis, and uncertainty – V: The discrete case. *Bell Syst Tech J* 57:1371–1430.
- Thompson DJ (1982) Spectrum estimation and harmonic analysis. *Proc IEEE* 70: 1055–1096.
- Nyquist H (1928) Certain topics in telegraph transmission theory. *Trans AIEE* 47: 617–644.
- Shannon CE (1937) Communication in the presence of noise. *Proc Inst Radio Eng* 37: 10–21.
- Kaulakys B, Gontis V, Alaburda M (2005) Point process model of 1/f noise vs a sum of lorentzians. *Phys Rev Lett* E71(5):1–11.